

WBJEE - 2017

Answer Keys by

Aakash Institute, Kolkata Centre

MATHEMATICS

Q.No.				
01	B	A	C	B
02	A	C	A	B
03	D	C	B	B
04	B	C	D	D
05	D	A	B	B
06	C	D	B	B
07	B	C	C	A
08	B	B	A	A
09	A	*	B	D
10	C	C	B	B
11	D	A	A	D
12	B	B	C	B
13	A	D	D	D
14	C	B	A	A
15	C	B	B	B
16	C	C	B	A
17	A	A	B	D
18	D	B	B	B
19	C	B	D	D
20	B	A	B	C
21	*	C	B	B
22	C	D	A	B
23	A	A	A	A
24	B	B	D	C
25	D	B	B	D
26	B	B	D	B
27	B	B	B	A
28	C	D	D	C
29	A	B	A	C
30	B	B	B	C
31	B	A	A	A
32	A	A	D	D
33	C	D	B	C
34	D	B	D	B
35	A	D	C	*
36	B	B	B	C
37	B	D	B	A
38	B	A	A	B
39	B	B	C	D
40	D	A	D	B
41	B	D	B	B
42	B	B	A	C
43	A	D	C	A
44	A	C	C	B
45	D	B	C	B
46	B	B	A	A
47	D	A	D	C
48	B	C	C	D
49	D	D	B	A
50	A	B	*	B
51	C	C	B	B
52	A	B	C	C
53	D	B	A	C
54	C	A	C	A
55	B	A	B	D
56	B	C	C	C
57	A	B	C	B
58	A	C	A	B
59	C	A	D	A
60	B	C	C	A
61	C	B	B	C
62	A	C	B	B
63	C	C	A	C
64	B	A	A	A
65	C	D	C	C
66	B	C	B,C	C,D
67	B,C	B,D	A,C	A,C
68	C	B,C	C	C
69	B,D	A,C	C,D	B
70	B,C	C	A,C	B,C
71	A,C	C,D	C	C
72	C	A,C	B	B,D
73	C,D	C	B,C	B,C
74	A,C	B	C	A,C
75	C	B,C	B,D	C

* Either B or D.



Aakash

Medical | IIT-JEE | Foundations

(Divisions of Aakash Educational Services Pvt. Ltd.)

Code - +

ANSWERS & HINT for WBJEE - 2017 SUB : MATHEMATICS

CATEGORY - I (Q1 to Q50)

Only one answer is correct. Correct answer will fetch full marks 1. Incorrect answer or any combination of more than one answer will fetch $-\frac{1}{4}$ marks. No answer will fetch 0 marks.

1. Transforming to parallel axes through a point (p, q) , the equation

$2x^2 + 3xy + 4y^2 + x + 18y + 25 = 0$ becomes $2x^2 + 3xy + 4y^2 = 1$. Then

- (A) $p = -2, q = 3$ (B) $p = 2, q = -3$ (C) $p = 3, q = -4$ (D) $p = -4, q = 3$

Ans : (B)

Hint : $4p + 3q + 1 = 0$

$3p + 8q + 18 = 0$

$\therefore p = 2, q = -3$

2. Let $A(2, -3)$ and $B(-2, 1)$ be two angular points of $\triangle ABC$. If the centroid of the triangle moves on the line $2x + 3y = 1$, then the locus of the angular point C is given by

- (A) $2x + 3y = 9$ (B) $2x - 3y = 9$ (C) $3x + 2y = 5$ (D) $3x - 2y = 3$

Ans : (A)


Hint : $G\left(t, \frac{1-2t}{3}\right), \alpha = 3t$

$\beta = 3 - 2t, \therefore 2x + 3y = 9$

3. The point $P(3, 6)$ is first reflected on the line $y = x$ and then the image point Q is again reflected on the line $y = -x$ to get the image point Q' . Then the circumcentre of the $\triangle PQQ'$ is

- (A) $(6, 3)$ (B) $(6, -3)$ (C) $(3, -6)$ (D) $(0, 0)$

Ans : (D)



4. Let d_1 and d_2 be the lengths of the perpendiculars drawn from any point of the line $7x - 9y + 10 = 0$ upon the lines $3x + 4y = 5$ and $12x + 5y = 7$ respectively. Then

- (A) $d_1 > d_2$ (B) $d_1 = d_2$ (C) $d_1 < d_2$ (D) $d_1 = 2d_2$

Ans : (B)

Hint :

5. The common chord of the circles $x^2 + y^2 - 4x - 4y = 0$ and $2x^2 + 2y^2 = 32$ subtends at the origin an angle equal to

- (A) $\frac{\pi}{3}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{6}$ (D) $\frac{\pi}{2}$

Ans : (D)


Hint : This common chord is passing through the centre of the 1st circle. Therefore it will form an angle of 90° at the circumferential point $(0, 0)$.

6. The locus of the mid-points of the chords of the circle $x^2 + y^2 + 2x - 2y - 2 = 0$ which make an angle of 90° at the centre is

- (A) $x^2 + y^2 - 2x - 2y = 0$ (B) $x^2 + y^2 - 2x + 2y = 0$ (C) $x^2 + y^2 + 2x - 2y = 0$ (D) $x^2 + y^2 + 2x - 2y - 1 = 0$

Ans : (C)

$$\sin 45^\circ = \frac{OP}{2} \Rightarrow OP = \sqrt{2}, \text{ Centre : } (-1, 1)$$




Hint :



7. Let P be the foot of the perpendicular from focus S of hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ on the line $bx - ay = 0$ and let C be the centre of hyperbola. Then the area of the rectangle whose sides are equal to that of SP and CP is

- (A) $2ab$ (B) ab (C) $\frac{(a^2 + b^2)}{2}$ (D) $\frac{a}{b}$

Ans : (B)




Hint : Area = $SP \cdot CP = a \cdot b$

8. B is an extremity of the minor axis of an ellipse whose foci are S and S'. If $\angle SBS'$ is a right angle, then the eccentricity of the ellipse is

- (A) $\frac{1}{2}$ (B) $\frac{1}{\sqrt{2}}$ (C) $\frac{2}{3}$ (D) $\frac{1}{3}$

Ans : (B)



$$b = ae, e = \frac{1}{\sqrt{2}}$$

9. The axis of the parabola $x^2 + 2xy + y^2 - 5x + 5y - 5 = 0$ is

- (A) $x + y = 0$ (B) $x + y - 1 = 0$ (C) $x - y + 1 = 0$ (D) $x - y = \frac{1}{\sqrt{2}}$

Ans : (A)

Hint : $(x + y)^2 = 5x - 5y + 5 \Rightarrow (x + y)^2 = 5(x - y + 1)$

\therefore Axis is $x + y = 0$

10. The line segment joining the foci of the hyperbola $x^2 - y^2 + 1 = 0$ is one of the diameters of a circle. The equation of the circle is

- (A) $x^2 + y^2 = 4$ (B) $x^2 + y^2 = \sqrt{2}$ (C) $x^2 + y^2 = 2$ (D) $x^2 + y^2 = 2\sqrt{2}$

Ans : (C)

Hint : $\therefore x^2 - y^2 + 1 = 0$, Foci = $(0, \pm\sqrt{2})$, Centre = $(0, 0)$, Radius = $\sqrt{2}$

Equation of circle $x^2 + y^2 = 2$

11. The equation of the plane through $(1, 2, -3)$ and $(2, -2, 1)$ and parallel to X-axis is

- (A) $y - z + 1 = 0$ (B) $y - z - 1 = 0$ (C) $y + z - 1 = 0$ (D) $y + z + 1 = 0$

Ans : (D)

Hint : $\begin{vmatrix} x-1 & y-2 & z+3 \\ 2-1 & -2-2 & 1+3 \\ 1 & 0 & 0 \end{vmatrix} = 0 \Rightarrow y+z+1=0$

12. Three lines are drawn from the origin O with direction cosines proportional to $(1, -1, 1)$, $(2-3, 0)$ and $(1, 0, 3)$. The three lines are

- (A) not coplanar (B) coplanar
(C) perpendicular to each other (D) coincident

Ans : (B)

Hint : $\Delta = 0$ (Coplanar)

13. Consider the non-constant differentiable function f of one variable which obeys the relation $\frac{f(x)}{f(y)} = f(x-y)$. If $f'(0) = p$ and $f'(5) = q$, then $f'(-5)$ is

(A) $\frac{p^2}{q}$ (B) $\frac{q}{p}$ (C) $\frac{p}{q}$ (D) q

Ans : (A)

Hint : $f(x) = a^{kx} \Rightarrow f'(x) = ka^{kx} \ln a$
 $k \ln a = p, ka^{5k} \ln a = q$

$$\Rightarrow a^{5k} = \frac{q}{p}$$

$$\therefore f'(-5) = k \cdot a^{-5k} \ln a = \frac{p^2}{q}$$

14. If $f(x) = \log_5 \log_3 x$, then $f'(e)$ is equal to

(A) $e \log_5 5$ (B) $e \log_5 3$ (C) $\frac{1}{e \log_e 5}$ (D) $\frac{1}{e \log_e 3}$

Ans : (C)

Hint : $f(x) = \log_5 \ln x + \log_5 \log_3 e$

$$f'(x) = \frac{1}{x} \cdot \frac{1}{\ln 5} \cdot \frac{1}{\ln x}$$

$$\therefore f'(e) = \frac{1}{e \ln 5}$$



15. Let $F(x) = e^x$, $G(x) = e^{-x}$ and $H(x) = G(F(x))$, where x is a real variable. Then $\frac{dH}{dx}$ at $x = 0$ is

(A) 1 (B) -1 (C) $-\frac{1}{e}$ (D) $-e$

Ans : (C)

Hint : $H(x) = e^{-e^x}$

$$\therefore H'(x) = -e^{-e^x} \cdot e^x$$

$$H'(0) = -\frac{1}{e}$$

16. If $f''(0) = k$, $k \neq 0$, then the value of $\lim_{x \rightarrow 0} \frac{2f(x) - 3f(2x) + f(4x)}{x^2}$ is

(A) k (B) $2k$ (C) $3k$ (D) $4k$

Ans : (C)

Hint : By L Hospital Rule

17. If $y = e^{m \sin^{-1} x}$, then $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - ky = 0$, where k is equal to

- (A) m^2 (B) 2 (C) -1 (D) $-m^2$

Ans : (A)

18. The chord of the curve $y = x^2 + 2ax + b$, joining the points where $x = \alpha$ and $x = \beta$, is parallel to the tangent to the curve at abscissa $x =$

- (A) $\frac{a+b}{2}$ (B) $\frac{2a+b}{3}$ (C) $\frac{2\alpha+\beta}{3}$ (D) $\frac{\alpha+\beta}{2}$

Ans : (D)

Hint : $2x+2a = (\beta+\alpha) + 2a$

$$\Rightarrow x = \frac{\alpha + \beta}{2}$$

19. Let $f(x) = x^{13} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 19$. Then $f(x) = 0$ has

- (A) 13 real roots (B) only one positive and only two negative real roots
(C) not more than one real root (D) has two positive and one negative real root

Ans : (C)

Hint : $f'(x) = 0$ has no real root

20. Let $f(x) = \begin{cases} \frac{x^p}{(\sin x)^q}, & \text{if } 0 < x \leq \frac{\pi}{2} \\ 0, & \text{if } x = 0 \end{cases}$, ($p, q, \in \mathbb{R}$). Then Lagrange's mean value theorem is applicable to $f(x)$ in closed

interval $[0, x]$

- (A) for all p, q (B) only when $p > q$ (C) only when $p < q$ (D) for no value of p, q

Ans : (B)

Hint : $\lim_{x \rightarrow 0^+} f(x) = 0$

$$\Rightarrow p > q$$

21. $\lim_{x \rightarrow 0} (\sin x)^{2 \tan x}$

- (A) is 2 (B) is 1 (C) is 0 (D) does not exist

Ans : Either B Or D

Hint : $\lim_{n \rightarrow 0^-} (\sin x)^{2 \tan x} \rightarrow$ Not in the domain hence does not exist, But if approached like

$$\lim_{n \rightarrow 0^-} (\sin^2 x)^{\tan x} = \lim_{n \rightarrow 0^+} (\sin^2 x)^{\tan x} = 1$$

22. $\int \cos(\log x) dx = F(x) + c$, where c is an arbitrary constant. Here $F(x) =$

- (A) $x[\cos(\log x) + \sin(\log x)]$ (B) $x[\cos(\log x) - \sin(\log x)]$
(C) $\frac{x}{2}[\cos(\log x) + \sin(\log x)]$ (D) $\frac{x}{2}[\cos(\log x) - \sin(\log x)]$

Ans : (C)

Hint : $\int \cos(\log x) dx = F(x) + c$, Let $\log x = t$, $I = \int e^t \cos t dt = e^t \cos t + e^t \sin t - I$,

$$\therefore I = \frac{e^t \cos t + e^t \sin t}{2} = \frac{x}{2} [\cos(\log x) + \sin(\log x)]$$

23. $\int \frac{x^2 - 1}{x^4 + 3x^2 + 1} dx (x > 0)$ is

- (A) $\tan^{-1}\left(x + \frac{1}{x}\right) + c$ (B) $\tan^{-1}\left(x - \frac{1}{x}\right) + c$ (C) $\log_e \left(\frac{x + \frac{1}{x} - 1}{x + \frac{1}{x} + 1} \right) + c$ (D) $\log_e \left(\frac{x - \frac{1}{x} - 1}{x - \frac{1}{x} + 1} \right) + c$

Ans : (A)

Hint : dividing by x^2 , $\int \frac{1 - 1/x^2}{x^2 + 1/x^2 + 3} dx$, Let $x + \frac{1}{x} = t$, $\int \frac{dt}{t^2 + 1} = \tan^{-1}\left(x + \frac{1}{x}\right) + c$

24. Let $I = \int_{10}^{19} \frac{\sin x}{1+x^8} dx$. Then

- (A) $|I| < 10^{-9}$ (B) $|I| < 10^{-7}$ (C) $|I| < 10^{-5}$ (D) $|I| > 10^{-7}$

Ans : (B)

Hint : $\left| \int_{10}^{19} \frac{\sin x}{1+x^8} dx \right| \leq \int_{10}^{19} \frac{|\sin x|}{1+x^8} dx \leq \int_{10}^{19} \frac{1}{1+x^8} dx$ (as $|\sin x| \leq 1$) $< \int_{10}^{19} 10^{-8} dx$ (as $1+x^8 > 10^8$ for $10 \leq x \leq 19$) $= 9 \times 10^{-8} < 10^{-7}$

25. Let $I_1 = \int_0^n [x] dx$ and $I_2 = \int_0^n \{x\} dx$, where $[x]$ and $\{x\}$ are integral and fractional parts of x and $n \in \mathbb{N} - \{1\}$. Then I_1/I_2 is equal to

- (A) $\frac{1}{n-1}$ (B) $\frac{1}{n}$ (C) n (D) $n-1$

Ans : (D)

Hint : $I_1 = \int_0^n [x] dx = \int_0^1 0 dx + \int_1^2 1 dx + \int_2^3 2 dx + \dots + \int_{n-1}^n (n-1) dx$, $= 0 + 1 + 2 + 3 + \dots + (n-1) = \frac{n(n-1)}{2}$,

$$I_2 = \int_0^n \{x\} dx = \int_0^n x dx - I_1 = \frac{n^2}{2} - \frac{n(n-1)}{2} = \frac{n}{2}, \therefore I_1/I_2 = n-1$$

26. The value of $\lim_{n \rightarrow \infty} \left[\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \dots + \frac{1}{2n} \right]$ is

- (A) $\frac{n\pi}{4}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{4n}$ (D) $\frac{\pi}{2n}$

Ans : (B)

Hint : $\lim_{n \rightarrow \infty} \left[\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \dots + \frac{1}{2n} \right] = \lim_{n \rightarrow \infty} \left[\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \dots + \frac{n}{n^2 + n^2} \right] = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \frac{1}{1 + \left(\frac{r}{n}\right)^2} = \int_0^1 \frac{1}{1+x^2} dx = \pi/4$

27. The value of the integral $\int_0^1 e^{x^2} dx$

- (A) is less than 1
 (B) is greater than 1
 (C) is less than or equal to 1
 (D) lies in the closed interval $[1, e]$

Ans : (B)

Hint : $\int_0^1 e^{x^2} dx > 1$

28. $\int_0^{100} e^{x-[x]} dx =$

- (A) $\frac{e^{100} - 1}{100}$ (B) $\frac{e^{100} - 1}{e - 1}$ (C) $100(e-1)$ (D) $\frac{e - 1}{100}$

Ans : (C)

Hint : $\int_0^{100} e^{x-[x]} dx$

$$= \int_0^1 e^x dx + \int_1^2 e^{x-1} dx + \int_2^3 e^{x-2} dx + \dots + \int_{99}^{100} e^{x-99} dx$$

$$= \int_0^1 e^x dx + \int_0^1 e^x dx + \int_0^1 e^x dx + \dots + \int_0^1 e^x dx$$

$$= 100 \times (e-1)$$

29. Solution of $(x + y)^2 \frac{dy}{dx} = a^2$ ('a' being a constant) is

- (A) $\frac{(x + y)}{a} = \tan \frac{y + c}{a}$, c is an arbitrary constant (B) $xy = a \tan cx$, c is an arbitrary constant
 (C) $\frac{x}{a} = \tan \frac{y}{c}$, c is an arbitrary constant (D) $xy = \tan(x+c)$, c is an arbitrary constant

Ans : (A)

Hint : $(x + y)^2 \frac{dy}{dx} = a^2$

[Put $x + y = z \Rightarrow 1 + \frac{dy}{dx} = \frac{dz}{dx}$]

$$\Rightarrow z^2 \left(\frac{dz}{dx} - 1 \right) = a^2 \Rightarrow z^2 + a^2 = z^2 \frac{dz}{dx}$$

$$\Rightarrow \int \frac{z^2}{z^2 + a^2} dz = \int dx \Rightarrow z - a \tan^{-1} \frac{z}{a} = x + C$$

$$\Rightarrow \frac{x + y}{a} = \tan \frac{y + c}{a}, \text{ c is arbitrary constant}$$

30. The integrating factor of the first order differential equation

$$x^2(x^2 - 1)\frac{dy}{dx} + x(x^2 + 1)y = x^2 - 1 \text{ is}$$

- (A) e^x (B) $x - \frac{1}{x}$ (C) $x + \frac{1}{x}$ (D) $\frac{1}{x^2}$

Ans : (B)

Hint : $x^2(x^2 - 1)\frac{dy}{dx} + x(x^2 + 1)y = x^2 - 1$, I.F = $e^{\int \frac{x(x^2+1)}{x^2(x^2-1)} dx}$,

$$e^{\int \frac{x^2-1}{x(x^2-1)} + \frac{2}{x(x+1)(x-1)} dx} = e^{\int \frac{1}{x} + \frac{-2}{x} + \frac{1}{x+1} + \frac{1}{x-1} dx} = e^{\ln\left(\frac{x^2-1}{x}\right)} = x - 1/x$$

31. In a G.P. series consisting of positive terms, each term is equal to the sum of next two terms. Then the common ratio of this G.P. series is

- (A) $\sqrt{5}$ (B) $\frac{\sqrt{5}-1}{2}$ (C) $\frac{\sqrt{5}}{2}$ (D) $\frac{\sqrt{5}+1}{2}$

Ans : (B)

Hint : $t_r = t_{r+1} + t_{r+2}$ $ar^{n-1} = ar^n + ar^{n+1} \Rightarrow 1 = r + r^2 \Rightarrow r = \frac{\sqrt{5}-1}{2}$

32. If $(\log_5 x)(\log_x 3x)(\log_{3x} y) = \log_x x^3$, then y equals

- (A) 125 (B) 25 (C) 5/3 (D) 243

Ans : (A)

Hint : $\frac{\log x \cdot \log 3x \cdot \log y}{\log 5 \cdot \log x \cdot \log 3x} = 3 \Rightarrow \log y = 3\log 5 \Rightarrow y = 5^3 = 125$

33. The expression $\frac{(1+i)^n}{(1-i)^{n-2}}$ equals

- (A) $-i^{n+1}$ (B) i^{n+1} (C) $-2i^{n+1}$ (D) 1

Ans : (C)

Hint : $(1-i) = \frac{2}{(1+i)}$

$$\frac{(1+i)^n}{(1-i)^{n-2}} = \frac{(1+i)^n}{2^{n-2}} \cdot (1+i)^{n-2} = \frac{(1+i)^{2(n-1)}}{2^{n-2}} = \frac{(2i)^{n-1}}{2^{n-2}} = 2i^{n-1} = -2i^{n+1}$$

34. Let $z = x + iy$, where x and y are real. The points (x, y) in the X-Y plane for which $\frac{z+i}{z-i}$ is purely imaginary lie on

- (A) a straight line (B) an ellipse (C) a hyperbola (D) a circle

Ans : (D)

Hint : Let $z = x + iy$

$$\therefore \frac{z+i}{(z-i)} = \frac{(x+i(y+1))(x+i(y-1))}{(x-i(1-y))(x+i(1-y))}$$

$$\operatorname{Re}\left(\frac{z+i}{z-i}\right) = 0 \Rightarrow \frac{x^2 + (y^2 - 1)}{x^2 + (1-y)^2} = 0 \Rightarrow x^2 + y^2 = 1$$

35. If p, q are odd integers, then the roots of the equation $2px^2 + (2p + q)x + q = 0$ are
 (A) rational (B) irrational (C) non-real (D) equal

Ans : (A)

Hint : $D = (2p + q)^2 - 8pq = (2p - q)^2 \rightarrow$ always a perfect square

36. Out of 7 consonants and 4 vowels, words are formed each having 3 consonants and 2 vowels. The number of such words that can be formed is

- (A) 210 (B) 25200 (C) 2520 (D) 302400

Ans : (B)

Hint : ${}^7C_3 \times {}^4C_2 \times 5! = 25200$

37. The number of all numbers having 5 digits, with distinct digits is

- (A) 99999 (B) $9 \times {}^9P_4$ (C) ${}^{10}P_5$ (D) 9P_4

Ans : (B)

Hint : $9 \times {}^9P_4$

38. The greatest integer which divides $(p + 1)(p + 2)(p + 3)\dots(p + q)$ for all $p \in \mathbb{N}$ and fixed $q \in \mathbb{N}$ is

- (A) $p!$ (B) $q!$ (C) p (D) q

Ans : (B)

Hint : This is product of ' q ' consecutive natural numbers, so it will always be divisible by $q!$

39. Let $((1 + x) + x^2)^9 = a_0 + a_1x + a_2x^2 + \dots + a_{18}x^{18}$. Then

- (A) $a_0 + a_2 + \dots + a_{18} = a_1 + a_3 + \dots + a_{17}$
 (B) $a_0 + a_2 + \dots + a_{18}$ is even
 (C) $a_0 + a_2 + \dots + a_{18}$ is divisible by 9
 (D) $a_0 + a_2 + \dots + a_{18}$ is divisible by 3 but not by 9

Ans : (B)

Hint : $a_0 + a_2 + a_4 + \dots + a_{18} = \frac{3^9 + 1}{2} \rightarrow$ even

40. The linear system of equations $\left. \begin{array}{l} 8x - 3y - 5z = 0 \\ 5x - 8y + 3z = 0 \\ 3x + 5y - 8z = 0 \end{array} \right\}$ has

- (A) only 'zero solution'
 (B) only finite number of non-zero solutions
 (C) no non-zero solution
 (D) infinitely many non-zero solutions

Ans : (D)

Hint : $D = \begin{vmatrix} 8 & -3 & -5 \\ 5 & -8 & 3 \\ 3 & 5 & -8 \end{vmatrix} = 0$

$D_1 = D_2 = D_3 = 0$ infinite solutions

41. Let P be the set of all non-singular matrices of order 3 over \mathbb{R} and Q be the set of all orthogonal matrices of order 3 over \mathbb{R} . Then,

- (A) P is proper subset of Q
- (B) Q is proper subset of P
- (C) Neither P is proper subset of Q nor Q is proper subset of P
- (D) $P \cap Q = \phi$, the void set

Ans : (B)

Hint : Q is the proper subset of P

42. Let $A = \begin{pmatrix} x+2 & 3x \\ 3 & x+2 \end{pmatrix}$, $B = \begin{pmatrix} x & 0 \\ 5 & x+2 \end{pmatrix}$. Then all solutions of the equation $\det(AB) = 0$ is

- (A) 1, -1, 0, 2
- (B) 1, 4, 0, -2
- (C) 1, -1, 4, 3
- (D) -1, 4, 0, 3

Ans : (B)

Hint : $\det|AB| = \begin{vmatrix} x^2+17x & 3x^2+6x \\ 8x+10 & (x+2)^2 \end{vmatrix} = 0$

$$\Rightarrow x(x+2)(x-4)(x-1) = 0$$

$$\Rightarrow x = 0, -2, 1, 4$$

43. The value of $\det A$, where $A = \begin{pmatrix} 1 & \cos\theta & 0 \\ -\cos\theta & 1 & \cos\theta \\ -1 & -\cos\theta & 1 \end{pmatrix}$ lies

- (A) in the closed interval [1, 2]
- (B) in the closed interval [0, 1]
- (C) in the open interval (0, 1)
- (D) in the open interval (1, 2)

Ans : (A)

Hint : $\det(A) = (1 + \cos^2\theta)$

$$\Rightarrow |A| \in [1, 2]$$

44. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be such that f is injective and $f(x) f(y) = f(x + y)$ for $\forall x, y \in \mathbb{R}$. If $f(x), f(y), f(z)$ are in G.P, then x, y, z, are in

- (A) A.P always
- (B) G.P always
- (C) A.P depending on the value of x, y, z
- (D) G.P depending on the value of x, y, z

Ans : (A)

Hint : $f(x) = a^x$

$$a^x, a^y, a^z \rightarrow \text{G.P}$$

$$x, y, z \rightarrow \text{A.P}$$

45. On the set \mathbb{R} of real numbers we define xPy if and only if $xy \geq 0$. Then the relation P is

- (A) reflexive but not symmetric
- (B) symmetric but not reflexive
- (C) transitive but not reflexive
- (D) reflexive and symmetric but not transitive

Ans : (D)

Hint : (-1, 0), (0, 2) satisfies the relation $xy \geq 0$ but (-1, 2) doesn't satisfy relation $xy \geq 0$.

46. On \mathbb{R} , the relation ρ be defined by 'xpy holds if and only if $x - y$ is zero or irrational'. Then

- (A) ρ is reflexive and transitive but not symmetric.
- (B) ρ is reflexive and symmetric but not transitive.
- (C) ρ is symmetric and transitive but not reflexive.
- (D) ρ is equivalence relation

Ans : (B)

47. Mean of n observations x_1, x_2, \dots, x_n is \bar{x} . If an observation x_q is replaced by x'_q then the new mean is

- (A) $\bar{x} - x_q + x'_q$
- (B) $\frac{(n-1)\bar{x} + x'_q}{n}$
- (C) $\frac{(n-1)\bar{x} - x_q}{n}$
- (D) $\frac{n\bar{x} - x_q + x'_q}{n}$

Ans : (D)

Hint : New Mean =
$$\frac{\sum_{i=1}^n x_i - x_q + x'_q}{n} = \frac{n\bar{x} - x_q + x'_q}{n}$$

48. The probability that a non leap year selected at random will have 53 Sundays is

- (A) 0
- (B) 1/7
- (C) 2/7
- (D) 3/7

Ans : (B)

49. The equation $\sin x (\sin x + \cos x) = k$ has real solutions, where k is a real number. Then

- (A) $0 \leq k \leq \frac{1+\sqrt{2}}{2}$
- (B) $2-\sqrt{3} \leq k \leq 2+\sqrt{3}$
- (C) $0 \leq k \leq 2-\sqrt{3}$
- (D) $\frac{1-\sqrt{2}}{2} \leq k \leq \frac{1+\sqrt{2}}{2}$

Ans : (D)

Hint : $\sin 2x - \cos 2x = 2k - 1$

$$\Rightarrow -\sqrt{2} \leq 2k - 1 \leq \sqrt{2}$$

$$\Rightarrow \frac{1-\sqrt{2}}{2} \leq k \leq \frac{\sqrt{2}+1}{2}$$

50. The possible values of x , which satisfy the trigonometric equation $\tan^{-1}\left(\frac{x-1}{x-2}\right) + \tan^{-1}\left(\frac{x+1}{x+2}\right) = \frac{\pi}{4}$ are

- (A) $\pm \frac{1}{\sqrt{2}}$
- (B) $\pm\sqrt{2}$
- (C) $\pm \frac{1}{2}$
- (D) ± 2

Ans : (A)

Hint :
$$\frac{x-1}{x-2} + \frac{x+1}{x+2} = 1 - \frac{x-1}{x-2} \cdot \frac{x+1}{x+2}$$

$$\Rightarrow x^2 + x - 2 + x^2 - x - 2 = x^2 - 4 - x^2 + 1$$

$$\Rightarrow 2x^2 = 1 \Rightarrow x = \pm \frac{1}{\sqrt{2}}$$

CATEGORY - II (Q51 to Q65)

Only one answer is correct. Correct answer will fetch full marks 2. Incorrect answer or any combination of more than one answer will fetch $-\frac{1}{2}$ marks. No answer will fetch 0 marks.

51. On set $A = \{1, 2, 3\}$, relations R and S are given by
 $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1)\}$
 $S = \{(1, 1), (2, 2), (3, 3), (1, 3), (3, 1)\}$ Then
 (A) $R \cup S$ is an equivalence relation
 (B) $R \cup S$ is reflexive and transitive but not symmetric
 (C) $R \cup S$ is reflexive and symmetric but not transitive
 (D) $R \cup S$ is symmetric and transitive but not reflexive


Ans : (C)

Hint : $R \cup S = \{(1,1), (2,2), (3,3), (1,2), (2,1), (1,3), (3,1)\}$

52. If one of the diameters of the curve $x^2 + y^2 - 4x - 6y + 9 = 0$ is a chord of a circle with centre $(1, 1)$, the radius of this circle is

- (A) 3 (B) 2 (C) $\sqrt{2}$ (D) 1

Ans : (A)



$$\therefore r = \sqrt{5+4} = 3$$



53. Let $A(-1, 0)$ and $B(2, 0)$ be two points. A point M moves in the plane in such a way that $\angle MBA = 2\angle MAB$. Then the point M moves along

- (A) a straight line (B) a parabola (C) an ellipse (D) a hyperbola

Ans : (D)

Hint :

54. If $f(x) = \int_{-1}^x |t| dt$, then for any $x \geq 0$, $f(x)$ is equal to

- (A) $\frac{1}{2}(1-x^2)$ (B) $1-x^2$ (C) $\frac{1}{2}(1+x^2)$ (D) $1+x^2$

Ans : (C)

Hint : $f(x) = \int_{-1}^x |t| dt, x \geq 0 = \frac{1}{2} (1^2 + x^2)$

55. Let for all $x > 0, f(x) = \lim_{n \rightarrow \infty} n \left(x^{\frac{1}{n}} - 1 \right)$, then

- (A) $f(x) + f\left(\frac{1}{x}\right) = 1$ (B) $f(xy) = f(x) + f(y)$ (C) $f(xy) = x f(y) + y f(x)$ (D) $f(xy) = x f(x) + y f(y)$

Ans : (B)

Hint : $f(x) = \lim_{n \rightarrow \infty} \frac{x^{1/n} - 1}{1/n} = \log x$

56. Let $I = \int_0^{100\pi} \sqrt{1 - \cos 2x} \, dx$, then

- (A) $I = 0$ (B) $I = 200\sqrt{2}$ (C) $I = \pi\sqrt{2}$ (D) $I = 100$

Ans : (B)


Hint : $I = \int_0^{100\pi} \sqrt{1 - \cos 2x} \, dx = 100\sqrt{2} \int_0^{\pi} |\sin x| \, dx = 200\sqrt{2}$

57. The area of the figure bounded by the parabolas $x = -2y^2$ and $x = 1 - 3y^2$ is

- (A) $\frac{4}{3}$ square units (B) $\frac{2}{3}$ square units (C) $\frac{3}{7}$ square units (D) $\frac{6}{7}$ square units

Ans : (A)

Hint : Curves intersect at $(-2, \pm 1)$



Area = $2 \int_0^1 (1 - y^2) \, dy = 2 \left(1 - \frac{1}{3}\right) = \frac{4}{3}$

58. Tangents are drawn to the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$ at the ends of both latus rectum. The area of the quadrilateral so formed is

- (A) 27 sq. units (B) $\frac{13}{2}$ sq. units (C) $\frac{15}{4}$ sq. units (D) 45 sq. units

Ans : (A)

Hint : Equation of tangent in quadrilateral I : $\frac{2x}{9} + \frac{y}{3} = 1$

59. The value of K in order that $f(x) = \sin x - \cos x - Kx + 5$ decreases for all positive real values of x is given by

- (A) $K < 1$ (B) $K \geq 1$ (C) $K > \sqrt{2}$ (D) $K < \sqrt{2}$

Ans : (C)

Hint : $f'(x) = \cos x + \sin x - k (< 0)$

$\therefore k > \cos x + \sin x$

max. $(\cos x + \sin x) = \sqrt{2}$

$\therefore k > \sqrt{2}$

60. For any vector \vec{x} , the value of $(\vec{x} \times \hat{i})^2 + (\vec{x} \times \hat{j})^2 + (\vec{x} \times \hat{k})^2$ is equal to

- (A) $|\vec{x}|^2$ (B) $2|\vec{x}|^2$ (C) $3|\vec{x}|^2$ (D) $4|\vec{x}|^2$

Ans : (B)

$$\begin{aligned} \text{Hint : } &= |\vec{x}|^2 (\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma) \\ &= 2|\vec{x}|^2 \end{aligned}$$

61. If the sum of two unit vectors is a unit vector, then the magnitude of their difference is

- (A) $\sqrt{2}$ units (B) 2 units (C) $\sqrt{3}$ units (D) $\sqrt{5}$ units

Ans : (C)

$$\begin{aligned} \text{Hint : } |\vec{a} - \vec{b}| &= \sqrt{|\vec{a}|^2 + |\vec{b}|^2 - 2\vec{a} \cdot \vec{b}} \\ &= \sqrt{3} \end{aligned}$$

62. Let α and β be the roots of $x^2 + x + 1 = 0$. If n be positive integer, then $\alpha^n + \beta^n$ is

- (A) $2\cos\frac{2n\pi}{3}$ (B) $2\sin\frac{2n\pi}{3}$ (C) $2\cos\frac{n\pi}{3}$ (D) $2\sin\frac{n\pi}{3}$

Ans : (A)

$$\text{Hint : } e^{i\frac{2n\pi}{3}} + e^{-i\frac{2n\pi}{3}} = 2\cos\left(\frac{2n\pi}{3}\right)$$

63. For real x , the greatest value of $\frac{x^2 + 2x + 4}{2x^2 + 4x + 9}$ is

- (A) 1 (B) -1 (C) $\frac{1}{2}$ (D) $\frac{1}{4}$

Ans : (C)

$$\text{Hint : } y = \frac{x^2 + 2x + 4}{2x^2 + 4x + 9} \text{ or, } (2y-1)(7y-3) \leq 0 \text{ or, } \frac{3}{7} \leq y \leq \frac{1}{2}$$

64. Let $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$. Then for positive integer n , A^n is

- (A) $\begin{pmatrix} 1 & n & n^2 \\ 0 & n^2 & n \\ 0 & 0 & n \end{pmatrix}$ (B) $\begin{pmatrix} 1 & n & n\left(\frac{n+1}{2}\right) \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix}$ (C) $\begin{pmatrix} 1 & n^2 & n \\ 0 & n & n^2 \\ 0 & 0 & n^2 \end{pmatrix}$ (D) $\begin{pmatrix} 1 & n & 2n-1 \\ 0 & \frac{n+1}{2} & n^2 \\ 0 & 0 & \frac{n+1}{2} \end{pmatrix}$

Ans : (B)

$$\text{Hint : } A = B + I \Rightarrow A^n = \begin{pmatrix} 1 & n & \frac{n(n+1)}{2} \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix}$$

65. Let a, b, c be such that $b(a+c) \neq 0$

$$\text{If } \begin{vmatrix} a & a+1 & a-1 \\ -b & b+1 & b-1 \\ c & c-1 & c+1 \end{vmatrix} + \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ (-1)^{n+2}a & (-1)^{n+1}b & (-1)^n c \end{vmatrix} = 0, \text{ then the value of } n \text{ is}$$

- (A) any integer (B) zero (C) any even integer (D) any odd integer

Ans : (C)

$$\text{Hint : } |A| = |A^T| \text{ or, } \begin{vmatrix} a & a+1 & a-1 \\ -b & b+1 & b-1 \\ c & c-1 & c+1 \end{vmatrix} - \begin{vmatrix} (-1)^{n+2}a & a+1 & a-1 \\ (-1)^{n+1}b & b+1 & b-1 \\ (-1)^n c & c-1 & c+1 \end{vmatrix} = 0 \Rightarrow (n+2) \text{ is even or } n \text{ is even}$$

CATEGORY - III (Q66 to Q75)

One or more answer(s) is (are) correct. Correct answer(s) will fetch full marks 2. Any combination containing one or more incorrect answer will fetch 0 marks. Also no answer will fetch 0 marks. If all correct answers are not marked and also no incorrect answer is marked then score = $2 \times \text{number of correct answer marked} \div \text{actual number of correct answers}$.

66. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be twice continuously differentiable. Let $f(0) = f(1) = f'(0) = 0$. Then

- (A) $f''(x) \neq 0$ for all x (B) $f''(c) = 0$ for some $c \in \mathbb{R}$ (C) $f''(x) \neq 0$ if $x \neq 0$ (D) $f'(x) > 0$ for all x

Ans : (B)

Hint : Applying Rolle's theorem twice $f'(x) = 0$ for some $x \in [0, 1]$

67. If $f(x) = x^n$, n being a non-negative integer, then the values of n for which $f'(\alpha+\beta) = f'(\alpha) + f'(\beta)$ for all $\alpha, \beta > 0$ is

- (A) 1 (B) 2 (C) 0 (D) 5

Ans : (B, C)

68. Let f be a non-constant continuous function for all $x \geq 0$. Let f satisfy the relation $f(x) f(a-x) = 1$ for some $a \in \mathbb{R}^+$. Then

$$I = \int_0^a \frac{dx}{1+f(x)}$$
 is equal to

- (A) a (B) $\frac{a}{4}$ (C) $\frac{a}{2}$ (D) f(a)

Ans : (C)

$$\text{Hint : } I = \int_0^a \frac{dx}{1+f(x)} = \int_0^a \frac{dx}{1+f(a-x)} = \int_0^a \frac{f(x)dx}{1+f(x)} \Rightarrow 2I = \int_0^a dx \Rightarrow I = \frac{a}{2}$$

69. If the line $ax + by + c = 0$, $ab \neq 0$, is a tangent to the curve $xy = 1-2x$, then

- (A) $a > 0, b < 0$ (B) $a > 0, b > 0$ (C) $a < 0, b > 0$ (D) $a < 0, b < 0$

Ans : (B,D)

$$\text{Hint : } \frac{dy}{dx} < 0$$

70. Two particles move in the same straight line starting at the same moment from the same point in the same direction. The first moves with constant velocity u and the second starts from rest with constant acceleration f . Then

- (A) they will be at the greatest distance at the end of time $\frac{u}{2f}$ from the start
- (B) they will be at the greatest distance at the end of time $\frac{u}{f}$ from the start
- (C) their greatest distance is $\frac{u^2}{2f}$
- (D) their greatest distance is $\frac{u^2}{f}$


Ans : (B, C)

Hint : $S = ut - \frac{1}{2}ft^2$

71. The complex number z satisfying the equation $|z-i| = |z+1| = 1$ is

- (A) 0
- (B) $1 + i$
- (C) $-1 + i$
- (D) $1 - i$

Ans : (A,C)



72. On \mathbb{R} , the set of real numbers, a relation ρ is defined as 'apb' if and only if $1 + ab > 0$. Then

- (A) ρ is an equivalence relation
- (B) ρ is reflexive and transitive but not symmetric
- (C) ρ is reflexive and symmetric but not transitive
- (D) ρ is only symmetric

Ans : (C)

73. If $a, b \in \{1, 2, 3\}$ and the equation $ax^2 + bx + 1 = 0$ has real roots, then

- (A) $a > b$
- (B) $a \leq b$
- (C) number of possible ordered pairs (a, b) is 3
- (D) $a < b$

Ans : (C, D)

Hint : $(1, 2) (1, 3) (2, 3)$

74. If the tangent to $y^2 = 4ax$ at the point $(at^2, 2at)$ where $|t| > 1$ is a normal to $x^2 - y^2 = a^2$ at the point $(a \sec \theta, a \tan \theta)$, then

- (A) $t = -\operatorname{cosec} \theta$
- (B) $t = -\sec \theta$
- (C) $t = 2 \tan \theta$
- (D) $t = 2 \cot \theta$

Ans : (A, C)

Hint : $x - yt = -at^2$ or, $\frac{x}{a \sec \theta} + \frac{y}{a \tan \theta} = 2 \Rightarrow t = -\operatorname{cosec} \theta$ or $t = 2 \tan \theta$

75. The focus of the conic $x^2 - 6x + 4y + 1 = 0$ is

- (A) $(2, 3)$
- (B) $(3, 2)$
- (C) $(3, 1)$
- (D) $(1, 4)$

Ans : (C)

Hint : $(x-3)^2 = -4(y-2)$

